

Second Generation Catalytic Asymmetric Synthesis of Tamiflu: Allylic Substitution Route


Tsuyoshi Mita, Nobuhisa Fukuda, Francesc X. Roca, Motomu Kanai,* and Masakatsu Shibasaki*

Graduate School of Pharmaceutical Sciences, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan

mshibasa@mol.f.u-tokyo.ac.jp

Received October 31, 2006

ABSTRACT



Catalytic asymmetric synthesis of Tamiflu, an important antiinfluenza drug, was achieved. After the catalytic enantioselective desymmetrization of *meso*-aziridine **3** with TMSN_3 , using a Y catalyst (1 mol %) derived from ligand **2**, an allylic oxygen function and C1 unit on the C=C double bond were introduced through cyanophosphorylation of enone and allylic substitution with an oxygen nucleophile. This second generation route of Tamiflu is more practical than our previously reported route.

The antiinfluenza drug Tamiflu (**1**)¹ is extremely important for protecting humans against a potential future pandemic of otherwise lethal flu. Considering the amount of Tamiflu required worldwide, there is an urgent demand to improve the Tamiflu production process. Currently, three groups have reported the asymmetric synthesis of Tamiflu:² Roche's commercial route utilizing naturally occurring shikimic acid as a starting material,³ Corey's route using the catalytic asymmetric Diels–Alder reaction developed by his group,⁴ and a route using the catalytic desymmetrization of *meso*-aziridines that was developed by our group.⁵ In this Letter,

we report an alternative route that significantly improves upon our initial synthesis.

Our previous synthesis of **1** utilized azido amide **4**, obtained with high enantioselectivity via catalytic desymmetrization of a *meso*-aziridine **3** as the starting chiral building block.⁵ A yttrium complex derived from ligand **2** was used as the asymmetric catalyst. There were two main drawbacks to this route: (1) allylic oxidation to introduce oxygen functionality to a diamine-derivative **10** required a stoichiometric amount of a toxic selenium reagent and (2) to effect this allylic oxidation in a synthetically useful yield, desymmetrized **4** was converted to a C_2 symmetric 1,2-diamine derivative (**10**: $P^1, P^2 = \text{Boc}$), which resulted in cumbersome protection group shuffling at the later stages of synthesis. Thus, we planned to develop a more practical synthetic route to circumvent these two problems (Scheme 1).

(1) Kim, C. U.; Lew, W.; Williams, M. A.; Liu, H.; Zhang, L.; Swaminathan, S.; Bischofberger, N.; Chen, M. S.; Mendel, D. B.; Tai, C. Y.; Laver, W. G.; Stevens, R. C. *J. Am. Chem. Soc.* **1997**, *119*, 681.

(2) For a recent review of synthetic strategies of Tamiflu, see: Farina, V.; Brown, J. D. *Angew. Chem., Int. Ed.* **2006**, *45*, 7330.

(3) Abrecht, S.; Harrington, P.; Iding, H.; Karpf, M.; Trussardi, R.; Wirz, B.; Zutter, U. *Chimia* **2004**, *58*, 621.

(4) Yeung, Y.-Y.; Hong, S.; Corey, E. J. *J. Am. Chem. Soc.* **2006**, *128*, 6310.

(5) Fukuta, Y.; Mita, T.; Fukuda, N.; Kanai, M.; Shibasaki, M. *J. Am. Chem. Soc.* **2006**, *128*, 6312.

attempted under thermal conditions.^{6b,12} Unexpectedly, a cyclic carbamate was cleanly produced via intramolecular S_N2' allylic substitution. After re-protection of the nitrogen atom of the cyclic carbamate with a Boc group to afford **16**, the less sterically hindered carbamate was cleaved by a catalytic amount of Cs₂CO₃ in MeOH, producing the corresponding α-OH allylic alcohol. The stereochemistry of the allylic position was inverted through an oxidation–stereoselective reduction⁵ sequence, affording trans amido alcohol **17** containing the suitable stereo-requirement for the following aziridine formation via the Mitsunobu reaction.¹³ The Mitsunobu reaction proceeded in high yield. The resulting aziridine **18** was subjected to a ring-opening reaction with 3-pentanol in the presence of BF₃·OEt₂,¹⁴ and the successive cleavage of the *N*-Boc group followed by *N*-acetylation afforded **19** in 81% yield. Hydrolysis of the *N*-Cbz group and ethanolysis of the nitrile proceeded concomitantly in acidic ethanol. Basification and H₃PO₄ salt formation¹⁵ completed the synthesis of Tamiflu (**1**).

Although this preliminary synthetic route avoided allylic oxidation with the toxic selenium reagent, protection group manipulation was still cumbersome and the synthesis was lengthy due to undesired iodocyclization with the acetamide group at the initial stage of the synthesis. To address this problem, we investigated a modified route involving iodocyclization of azido amide **11** (Scheme 3). Treatment of **11** with I₂ in the presence of K₂CO₃, followed by subjecting the resulting cyclic carbamate to an elimination reaction of HI with DBU, produced **20** in 85% yield. Protection of the carbamate nitrogen atom with a Boc group and reductive acetylation of the azide with AcSH¹⁶ produced **21**, which was selectively hydrolyzed and oxidized with Dess–Martin periodinane to produce enone **22**. Cyanophosphorylation of enone **22** proceeded with DEPC in the presence of LiCN, and cyanophosphate **23** was obtained as a single detectable isomer.

The key allylic rearrangement of **23** was studied under thermal conditions (Scheme 4). Heating a toluene solution of **23** to 140 °C in a sealed tube produced a labile and unisolatable β-allyl phosphate **28** as an initial product, whose structure was estimated by ESI-MS without isolation.¹⁷ Addition of 1 M NaOH solution to the reaction mixture produced stable oxazoline **29** as a major product in 38% yield, together with allylic alcohol **24** (23%) produced through an S_N2 attack of a hydroxide anion to the phosphate. Because we could not find any conditions that affect the

(11) Configuration of the tetrasubstituted carbon of **15** was temporarily assigned as shown based on the structure of **29** (see Scheme 4).

(12) Pd(II)-catalyzed allylic rearrangement (ref 6a) did not afford the desired product.

(13) Ho, M.; Chung, J. K. K.; Tang, N. *Tetrahedron Lett.* **1993**, *34*, 6513.


(14) Ring-opening reaction of cyclic carbamate **16** with 3-pentanol was intensively studied in the presence of a variety of Lewis acids. The desired product, however, was not produced in these reactions.

(15) Harrington, P. J.; Brown, J. D.; Foderaro, T.; Hughes, R. C. *Org. Process Res. Dev.* **2004**, *8*, 86.

(16) Shangquan, N.; Katukojvala, S.; Greenberg, R.; Williams, L. J. *J. Am. Chem. Soc.* **2003**, *125*, 7754.


(17) The stereochemistry of the allylic position of **28** was temporarily assigned based on Bartlett and Kurihara's previous findings (ref 6), in which allylic rearrangement of α-cyanophosphate proceeded in a suprafacial fashion.

Scheme 3. Optimized Allylic Substitution Route



oxazoline opening of **29** with 3-pentanol, we next turned our attention to an intermolecular nucleophilic substitution of the allylic phosphate of **28**. After several trials,¹⁸ a synthetically useful yield (78%) of α-allylic alcohol **24** was produced by the addition of saturated aqueous NH₄Cl solution to **28**.

Scheme 4. Allylic Rearrangement–Substitution Sequence



On the basis of the above findings, asymmetric synthesis of Tamiflu was completed from **24** as follows (Scheme 3).

(18) Neither direct addition of 3-pentanol to **28** nor aziridine formation from **28** proceeded under various conditions.

After the α -allyl alcohol of **24** was inverted via the Mitsunobu reaction with *p*-nitrobenzoic acid followed by hydrolysis of the *p*-nitrobenzoate in one pot, the resulting β -allylic alcohol **25** was subjected to aziridine formation through Mitsunobu conditions, producing **26**. Introduction of 3-pentanol was conducted with an aziridine opening reaction to produce **27**⁵ in 56% yield. Ethanolysis of the cyanide and cleavage of the Boc group under acidic ethanol, followed by basification and H₃PO₄ treatment,¹⁵ produced **1**. This route solves the two main problems (allylic oxidation and protection group shuffling) of the previously reported synthetic route.⁵ In addition, the synthetic scheme is shorter than that of the previously reported synthesis⁵ (15 steps from aziridine **3**).

In summary, we developed our second generation synthesis of Tamiflu. This route utilizes a catalytic enantioselective desymmetrization of *meso*-aziridine **3** with TMSN₃ as the initial key step. The reaction was performed with 1 mol % of chiral Y catalyst derived from ligand **2** at room temper-

ature. Careful optimization of the synthetic scheme made it possible to avoid protection group shuffling, which was a drawback in our first generation synthesis.⁵ The allylic oxygen functionality required for 3-pentyl ether formation was introduced by rearrangement of an allylic phosphate followed by substitution with a hydroxyl group. Thus, the other drawback of our previous synthesis—i.e., the use of a toxic selenium reagent in the allylic oxidation—was also overcome. Further improvement of the synthetic efficacy by using a completely different strategy is currently under investigation.

Acknowledgment. Financial support was provided by a Grant-in-Aid for Specially Promoted Research of MEXT.

Supporting Information Available: Experimental procedures and characterization of the products. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OL062663C