

Synthesis of (-)- Δ^9 -*trans*-Tetrahydrocannabinol: Stereocontrol via Mo-Catalyzed Asymmetric Allylic Alkylation Reaction


Barry M. Trost* and Kalindi Dogra

Department of Chemistry, Stanford University, Stanford, California 94305

bmtrost@stanford.edu

Received December 14, 2006

ABSTRACT



Δ^9 -THC is synthesized in enantiomerically pure form, where all of the stereochemistry is derived from the molybdenum-catalyzed asymmetric alkylation reaction of the extremely sterically congested bis-ortho-substituted cinnamyl carbonate in high regio- and enantioselectivity.

(-)- Δ^9 -*trans*-Tetrahydrocannabinol (Δ^9 -THC, **1**) isolated¹ from female *Cannabis sativa* L. in 1964 has been identified as the primary psychomimetic component of marijuana. It is also known to show antiemetic, antiglaucoma, and analgesic properties. Currently it is administered as an antiemetic to patients undergoing chemotherapy. Discovery of the cannabinoid receptors CB1 and CB2 and Δ^9 -THC analogues² that bind selectively to them has led to a need for the development of a flexible synthetic route that would yield target compounds easily in high yields and in stereochemically pure form.

THC itself has been prepared numerous times, though most routes are either racemic or derive their chirality from chiral building blocks.³ Only Evan's⁴ route targets the natural

product enantioselectively from achiral starting materials. The problems associated with the synthesis of THC involve the control of cis–trans stereochemistry at the cyclohexene ring and the position of the double bond, i.e., Δ^9 vs Δ^8 , the latter being thermodynamically more stable.

Our retrosynthetic analysis of the molecule envisioned all of the stereochemistry resulting from a single Mo-catalyzed asymmetric allylic alkylation (AAA) reaction (see the abstract). Alkylation of malonate adduct **4** with **3** should


(3) Selected examples: (a) Mechoulam, R.; Gaoni, Y. *J. Am. Chem. Soc.* **1965**, *87*, 3273. Mechoulam, R.; Braun, P.; Gaoni, Y. *J. Am. Chem. Soc.* **1967**, *89*, 4552. Mechoulam, R.; Braun, P.; Gaoni, Y. *J. Am. Chem. Soc.* **1972**, *94*, 6159. (b) Fahrenholtz, K. E.; Lurk, M.; Kierstead, R. W. *J. Am. Chem. Soc.* **1966**, *88*, 2079. Fahrenholtz, K. E.; Lurk, M.; Kierstead, R. W. *J. Am. Chem. Soc.* **1967**, *89*, 5934. (c) Chan, T. H.; Chaly, T. *Tetrahedron Lett.* **1982**, *23*, 2935. (d) Rickards, R. W.; Ronneberg, H. *J. Org. Chem.* **1984**, *49*, 5276. (e) Childers, W. E., Jr.; Pinnick, H. W. *J. Org. Chem.* **1984**, *49*, 5276. (f) Malkov, A. V.; Koovsky, P. *Collect. Czech. Chem. Commun.* **2001**, *66* (8), 1257. (g) William, A. D.; Kobayashi, Y. *J. Org. Chem.* **2002**, *67*, 8771. William, A. D.; Kobayashi, Y. *Org. Lett.* **2001**, *3*, 2017.

(4) (a) Evans, D. A.; Shaughnessy, E. A.; Barnes, D. M. *Tetrahedron Lett.* **1997**, *38*, 3193. (b) Evans, D. A.; Barnes, D. M.; Johnson, J. S.; Lectka, T.; Matt, P. V.; Miller, S. J.; Murry, J. A.; Norcross, R. D.; Shaughnessy, E. A.; Campos, K. R. *J. Am. Chem. Soc.* **1999**, *121*, 7582.

(1) Mechoulam, R.; Gaoni, Y. *J. Am. Chem. Soc.* **1964**, *86*, 1646.
(2) (a) Mahadevan, A.; Siegel, C.; Martin, B. R.; Abood, M. E.; Belestskaya, I.; Razdan, R. K. *J. Med. Chem.* **2000**, *43*, 3778. (b) Drake, D. J.; Jensen, R. S.; Busch-Petersen, J.; Kawakami, J. K.; Fernandez-Garcia, M. C.; Fan, P.; Makriyannis, A.; Tius, M. A. *J. Med. Chem.* **1998**, *41*, 3596. (c) Harrington, P. E.; Stergiades, I. A.; Erickson, J.; Makriyannis, A.; Tius, M. A. *J. Org. Chem.* **2000**, *65*, 6576. (d) Tius, M. A.; Makriyannis, A.; Zou, X. L.; Abadji, V. *Tetrahedron* **1994**, *50*, 2671.


furnish the substrate for ring-closing metathesis (RCM). The RCM in itself results in a fixed geometry of double bond thus solving the problem of co-formation of Δ^8 -THC seen in many syntheses. Decarboxylation of the malonate after RCM was planned to result in the required trans stereochemistry of **2**. Grignard addition to ester **2** followed by demethylation to the free phenol and cyclization yields **1**. Varying the aromatic group in **5**, the alkylating partner **3**, or the Grignard reagents one can potentially prepare different analogues of THC without fundamentally changing the chemistry involved.

Allyl alcohol **9** was easily prepared in high yield from commercially available olivetol, **6**, in 4 steps (Scheme 1).



Lithiated dimethyl olvitol was quenched with dry DMF to yield the aldehyde **8** in 83% yield. Wadsworth–Emmons reaction of **8** with sodium triethylphosphonoacetate resulted in the corresponding α,β -unsaturated ethyl ester, which was subjected without further purification to DIBAL-H reduction to yield 97% of **9**. Preparation of carbonate **5** proved to be a little tricky because the compound is sensitive to both acid and base, including silica chromatography conditions. A preparation of **5** clean enough to take into the Mo-alkylation reaction was achieved by titrating alcohol **9** with BuLi at $-78\text{ }^{\circ}\text{C}$ in ether and quenching the resulting alkoxide with methyl chloroformate also at $-78\text{ }^{\circ}\text{C}$. Washing the organic layer with ice-cold water followed by solvent removal gave carbonate **5** as a waxy solid, which is stable to storage.

The branched allyl alcohol **10** is prepared by condensation of acrolein with lithiated **7** in one step in 91% yield (Scheme 2). We planned to use the carbonate prepared from **10** in



the Mo-AAA reaction to reduce the number of steps and increase the efficiency of the synthesis. Unfortunately it proved impossible to prepare either the carbonate or the acetate of **10**, due to their facile decomposition aided by the highly electron rich aromatic ring.

We did, however, take advantage of this to isomerize the branched alcohol **10** into the linear alcohol **9**.⁵ Treatment of

the lithium alkoxide of **10** with dry CO_2 gas gives the corresponding carbonate, which rearranges in the presence of $\text{PdCl}_2(\text{CH}_3\text{CN})_2$ to yield the more stable linear carbonate that decomposes to the alcohol **9** on workup (Figure 1). A





Figure 1. Isomerization of branched **10** to linear alcohol **9**.

74% yield of **9** was observed with 6% recovered **10**. To the best of our knowledge this reaction has not been previously reported. This alternate strategy using this novel isomerization should prove to be a more efficient, atom economical approach to cinnamyl alcohols than the more traditional protocol based upon olefination.

With the carbonate in hand we began examining the allylic alkylation reaction. Although molybdenum is known for giving the product resulting from attack at the more substituted carbon, we were concerned that the two *o*-methoxy groups on the aryl ring would make this reaction sterically unfavorable. Nevertheless, the reaction of carbonate **5** with sodium dimethyl malonate under standard conditions of 10 mol % $[\text{Mo}(\text{CO})_3\text{C}_7\text{H}_8]$ and 15 mol % chiral ligand *S,S*-**11** proceeded sluggishly but gratifyingly gave only the branched product **4** in 95% yield and 95% enantiomeric excess. The reaction was optimized to reduce catalyst loading and the optimized conditions are shown in eq 1.




Attempts to alkylate the malonate adduct **4** with **3** under numerous conditions met with no success (Scheme 3). The electrophile **3** slowly disappeared under the reaction conditions due to 1,2-elimination, giving rise to isoprene. To examine whether this or the steric demands of making a quaternary center at the congested malonate carbon was the cause for failure, we tried the addition of MeI, but observed only trace amounts of the desired adduct. This led us to believe steric congestion was the cause of failure, and hence we prepared the monoester **12** under Krapcho decarboxylation⁶ conditions in 83% yield, which reduced steric demands. Since the enolates of methyl esters are generally stable only below $-35\text{ }^{\circ}\text{C}$ we attempted to carry out the alkylation at

(5) (a) Overman, L. E. *J. Am. Chem. Soc.* **1974**, 96, 597. (b) Overman, L. E. *Angew. Chem., Int. Ed. Engl.* **1984**, 23, 579.

(6) Krapcho, A. P.; Weimaster, J. F.; Eldridge, J. M.; Jahngen, G. E.; Lovey, A. J.; Stephens, W. P. *J. Org. Chem.* **1978**, 43, 138.


Scheme 3



–40 °C. Our best result was only a 50% yield of the adduct **13** using triflate **3c** (Scheme 3). The competing elimination reaction was still a problem due to the slower rate of alkylation at the reduced temperature.


Therefore a nucleophile whose enolate would be stable at room temperature or higher was required. We believed that the dianion of the acid **14** would meet our needs (Scheme 4). This acid is prepared from **4** using classical conditions

Scheme 4



in 97% yield, as shown in Scheme 5. The dianion of acid **14** cleanly underwent alkylation with iodide **3b** to yield 84% of **15** as a 2.4:1 mixture of anti and syn isomers at the newly formed center. The two isomers were separated using column

Scheme 5



chromatography, converted to their respective methyl esters, and subjected to ring closing metathesis using the second generation Grubb's ruthenium carbene catalyst,⁷ providing the anti and syn cyclohexene compounds **2** and **16**. The syn compound **16** was easily recycled by equilibrating to anti **2** by treatment with NaOMe in MeOH for 3 days. Addition of MeLi to the cyclized ester **2** at –78 °C resulted in formation of the tertiary alcohol **17** in 92% yield (Scheme 4).

With the end in sight, compound **17** was treated with BBr₃ to yield the free phenol by demethylation of both ethers (Scheme 5). We hoped that the free bis-phenol would subsequently cyclize under the acidic reaction conditions to give **1**. Unfortunately we got a complex mixture of products. The use of other acidic reagents for demethylation resulted in similar fates. Finally we decided to carry out the reaction in a stepwise manner using NaSEt in DMF, which is known to stop at monodeprotection of aryl diethers.⁸ The reaction proceeded as expected to yield 97% of **18**, [α]_D²³ –52 (*c* 0.79, CHCl₃) (lit.⁹ [α]_D²⁶ –44 (*c* 0.20, CHCl₃)). Formation of the cyclized ether was achieved by treating **18** with ZnBr₂ in the presence of MgSO₄. The resulting compound was subjected to NaSEt in DMF without further purification to yield the desired Δ^9 -THC **1** in 61% yield [α]_D²⁵ –152 (*c* 0.46, CHCl₃) (lit.⁶ [α]_D²⁸ –150 (*c* 1.0, CHCl₃)).

In summary, the use of the molybdenum-catalyzed asymmetric alkylation reaction developed in this group provided the stereochemical framework to synthesize Δ^9 -THC in enantiomerically pure form and 30% overall yield from olivetol dimethyl ether. The regio- and enantioselectivity of the Mo AAA is undiminished even in this sterically congested example. Furthermore, a simple and efficient two-step protocol for the synthesis of cinnamyl alcohols has also been developed.

Acknowledgment. We thank the General Medical Sciences Institute and the National Institutes of Health (GM 13598) and Merck for their generous support of our programs. We are also indebted to Johnson Matthey for a generous gift of palladium salts. Mass spectra were provided by the Mass Spectrometry Regional Center of the University of California, San Francisco supported by the NIH Division of Research Resources.

Supporting Information Available: Experimental procedures for the preparation of new compounds and characterization data. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OL063022K

(7) Scholl, M.; Ding, S.; Lee, C. W.; Grubbs, R. H. *Org. Lett.* **1999**, *1*, 953.

(8) Feutrill, G. I.; Mirrington, R. N. *Tetrahedron Lett.* **1970**, 1327.

(9) William, A. D.; Kobayashi, Y. *J. Org. Chem.* **2002**, *67*, 8771.